

CODEINE for Pain and Cough Relief \$@Buy Online with Fast UK Shipping AND (Affordable Rates)



What Is Codeine?

Codeine is an opiate and a parent drug for *morphine* that is mainly used to treat pain, diarrhea, and coughing. It is available as a single-ingredient product to relieve mild to moderate pain that requires opioid treatment.

It is often used as a combination medication with Acetaminophen, which helps relieve pain in different parts of the body, and it helps treat cough when combined with *NSAIDs such as Ibuprofen*.

Codeine is approved for the treatment of pain and cough by U.S. food and drug administration, but FDA restricts its use for childrens because it does not comply with its safe to use regulation.

According to the World Health Organization (WHO), Codeine is considered an essential element for adults. Medicines referring to this category are made for adult use only and not for children below *12 years of age*.

Codeine is a round-shaped white tablet that is available in various strengths such as *15mg, 30mg, 300/30mg, and 60 mg at 'United State Pharmacy' at up to 25% discount*.

Why Codeine Is Used

- Mild To Moderate Pain

Codeine helps in treating mild to moderate pain that requires opioid treatment, and other alternative therapies are ineffective and are easily tolerated.

- Diarrhea

It is recommended for short-term treatment in people suffering from diarrhea, as it helps in slowing down bowel movement that causes constipation.

- Cough

Codeine is also used to treat cough in adults, as it works by decreasing the activity in the part of the brain that causes coughing.

How It Works

It works on the central nervous system (CNS) by blocking the way the brain signals pain to the body to induce insensibility to pain. Codeine binds to the opioid receptors, promoting relief and instant relaxation from pain.

What Should I Tell My Healthcare Provider Before Using Codeine?

Anyone suffering from pain, diarrhea, and cough can use it safely on a doctor's prescription, but people with certain health conditions are not recommended to use it.

Tell your doctor before using codeine if you have –

- Gallstones
- Head injury
- Ulcerative colitis
- Enlarged prostate
- Myasthenia gravis
- Breastfeeding mother
- Adrenal gland problems
- Underactive thyroid gland
- Obstructive sleep apnea (OSA)
- Allergic to Codeine or its ingredients
- Suffering from kidney, liver, or lung problems

How To Take The Dose

For Adults –

- The recommended dose is 15 – 16 mg to be taken every 4 hours. Your doctor may adjust or increase your dose according to its response. Do not take more than a 360 mg dose in a day.

For children (12 – 18 years) –

- Children below 18 years of age should not use this medication without consulting a doctor.

Possible Side Effects Of Codeine

Codeine causes mild to moderate side effects when used in inadequate amounts or without a doctor's prescription.

Some of the common and most notable side effects of Codeine are –

- Headache
- Xerostomia
- Dry mouth
- Sick feeling
- Vomiting
- Constipation

Serious Side Effects

- Insomnia – Codeine may affect your sleep schedule by causing overthinking problems and worsening anxiety when used in inadequate amounts.
- Fast heartbeat – Your heart may start to beat faster or slower. In these conditions, talk to your doctor.
- Urinary retention – Codeine can cause urinary tract infections. Speak to your doctor if you experience frequent urination, urgency to urinate, pain or burning sensation during urination, or blood in the urine.
- Xerostomia – Dry, rough, or raw tongue, loss of taste, and sore throat are signs of Xerostomia (a medical condition).
- CNS depression – Overdose of Codeine can cause symptoms like headaches, loss of feeling, weakness, and changes in vision, hearing, and taste, which are signs of CNS depression.
- Low blood pressure – It can cause hypotension, and you may feel lightheaded, faint, or have blurred vision or fatigue.
- Low adrenal gland function – You may experience nausea, vomiting, loss of appetite, unusual weakness, or fatigue.
- Allergic Reactions – Rashes on the skin, itching, hives, swelling on the face, lips, throat, or tongue.

What To Prevent While Using Codeine

You must remember these preventative measures in order to prevent any severe side effects, drug interaction, and withdrawal symptoms.

- Tell your doctor if you have an underactive thyroid, enlarged prostate, or urinary problems before using it to prevent any harmful effects of this medication.
- Avoid using alcohol, other substances, or drugs while using this medication to prevent any drug interaction that can cause any serious withdrawal symptoms.
- Take the dose of Codeine with food or just after having a meal or snack to ease the feeling of sickness.
- Breastfeeding mothers should not use Codeine, as it can be harmful to the health of the baby and the mother and can cause respiratory issues or even death of the nursing infant.